

Improving your bladder control

There are other simple measures you can take to promote a healthy bladder and pelvic floor.

- To prevent or reduce urine leakage, squeeze the pelvic floor muscles tightly *at the same time* as you do any activity that usually causes urine leakage. This procedure is called *The Knack*
- Maintain a reasonable fluid intake for your weight of approximately 30ml per kg in 24 hours
- Reduce the intake of caffeinated drinks such as tea, coffee, chocolate and fizzy drinks, as these can increase bladder and bowel symptoms
- Avoid constipation and straining

- Avoid heavy lifting
- If overweight, aim to reduce your weight to an acceptable level for your height and build
- A persistent cough can make it more likely that you leak urine. It is very important that you always tighten your pelvic floor muscles during any coughing and also seek advice from your Doctor
- Smoking can make a cough worse and also increases the risk of cancers, including bladder cancer. If you would like help in stopping smoking, see your Doctor or Nurse for advice
- If you see any blood in your urine, you must contact your Doctor or Nurse for advice

The After-dribble

A very common and embarrassing problem is a small urine leak from the water pipe (urethra) after leaving the toilet.

This can result in a damp patch, especially noticeable on light coloured trousers.

This can be prevented by either tightening the pelvic floor muscles after fully emptying the bladder, or by placing your hand behind the scrotum and applying firm pressure upwards and slightly forward. Any urine left in the water pipe (urethra) will be expelled into the urinal or toilet.

DO YOU KNOW? 90% of men aged 50 to 80 years old suffer from bladder problems.¹

If the advice given in this leaflet does not improve your symptoms, further advice and help is available from:

- Your **GP**, who may also refer you to other Specialists for advice.
 - The **Practice Nurse, District Practitioner, District Nurse or Health Visitor** may be able to give further advice.
 - A **Continence Specialist or Advisor** is an experienced Specialist Nurse, Physiotherapist or other Health Care Professional.
 - A Urologist specialises in the branch of medicine dealing with the genitourinary system.
- The decision to refer you to the Urologist is usually made by your GP. In some areas, other professionals can make this referral.
- Details of the services available in your area are available from your Health Trust, Clinic or Health Centre, or GP Practice.**
- There are many useful organisations which may be of further help and support, such as the Bladder and Bowel Foundation at www.bladderandbowelfoundation.org Tel: 0845 345 0165
- Details of other organisations are available from your GP or local Continence Service.

1. NICE Clinical Guideline 97 (2010)

PAUL HARTMANN Limited
Heywood Distribution Park
Pilsworth Road
Heywood
OL10 2TT
Tel: 01706 363200
Fax: 01706 363201
www.hartmann-direct.co.uk



Going further
for health

HARTMANN



Going further
for health

Continence
Management

Pelvic Floor Muscle Exercises for Men

Improving your bladder control




About your pelvic floor muscles

The pelvic floor muscles extend from the pubic bone to the base of the spine (see diagram).

There are two openings in the pelvic floor, one for the urethra, sometimes called the 'water pipe' (the passageway that carries urine from the bladder), and one through which the bowel empties.

The pelvic floor muscles:

- Support the lower abdominal contents, especially the bladder and bowel
- Assist normal bladder and bowel function and prevent leakage (incontinence)
- Contribute to sexual function



What might affect the pelvic floor muscles?

In men, the pelvic floor muscles may be affected as a result of:

- Surgery on the prostate gland
- Persistent constipation
- Lifting heavy loads at work or home
- Chest problems with a chronic cough
- Being overweight
- Being generally unfit
- Some medical conditions

Weak pelvic floor muscles can result in urine leakage on effort or exertion, or on sneezing, coughing or laughing

How to strengthen your pelvic floor muscles

You will need to learn, and regularly carry out, your pelvic floor muscle training programme. To begin, choose a quiet time and place, so you can concentrate on learning the exercises correctly.

Sit, stand or lie down, with your feet comfortably apart. You can do the exercises in any of these positions. Relax and breathe normally.



Pelvic floor muscle exercises

You are going to learn to carry out both slow and fast pelvic floor muscle exercises.

☺ Concentrate, then squeeze and lift the pelvic floor muscles as if you are trying to stop yourself from passing urine. Then tighten the muscles around the back passage, as if you are trying not to pass wind. By doing these two together, you should be exercising your pelvic floor muscles. As you tighten the muscles, you will feel your scrotum and base of your penis move slightly upwards.

☺ **Slow exercise**
Tighten your pelvic floor muscles and squeeze for as long as you can, then 'let go'

and relax for about 4 seconds, then repeat. You may only be able to hold for a few seconds to begin with, but your aim is to hold for 10 seconds, repeated at least 8 times. To begin with, watch a second hand on a clock to time this.

☺ **Fast exercise**
'Squeeze and let go' your pelvic floor muscles in the same way as before but much faster – about one per second for 10 seconds. Your penis should twitch upwards with each contraction

☺ **You have the best chance of success if you practice doing these slow and fast contractions 3 times every day for at least three months.**

Each month, you can monitor your progress below:

Date	Hold for?	Number of slow squeezes	Number of fast squeezes
(Start)	secs	times	times
	secs	times	times
	secs	times	times
	secs	times	times
	secs	times	times